

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28 29 30 31 32 33 34 35 36 37 38 39 40 41 42 43 44 45 46 47 48 49 50 51 52 53 54 55 56 57 58 59 60 61 62 63 64 65 66 67 68 69 70 71 72 73 74 75 76 77 78 79 80 81 82 83 84 85 86 87 88 89 90 91 92 93 94 95 96 97 98 99 100

(43) International Publication Date
12 April 2001 (12.04.2001)

PCT

(10) International Publication Number
WO 01/25410 A2

(51) International Patent Classification⁷: C12N 9/72,
C07K 14/81, 5/00, C12Q 1/37, G01N 33/68, A61K 38/49,
38/04, 38/57

(74) Agent: LIVNAT, Shmuel; Venable, P.O. Box 34385,
Washington, DC 20043-9998 (US).

(21) International Application Number: PCT/US00/26502

(22) International Filing Date:
27 September 2000 (27.09.2000)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/157,012 1 October 1999 (01.10.1999) US

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ,
DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR,
HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR,
LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ,
NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM,
TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,
IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG,
CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).


(71) Applicant: ANGSTROM PHARMACEUTICALS,
INC. [US/US]; Suite 105, 6310 Nancy Ridge Drive, San
Diego, CA 92121 (US).

Published:
— Without international search report and to be republished
upon receipt of that report.

(72) Inventors: MAZAR, Andrew, P.; 8650-4 Villa La Jolla
Drive, La Jolla, CA 92037 (US). JONES, Terence, R.;
4915 Quincy Street, San Diego, CA 92109 (US).

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: DIAGNOSTIC PROBES AND THERAPEUTICS TARGETING uPA AND uPAR



Urokinase Plasminogen Activator (uPA)

WO 01/25410 A2

(57) Abstract: A uPAR-targeting protein or peptide is diagnostically or therapeutically labeled and used in methods of diagnosis of therapy. The labeled protein or peptide preferably has the following properties: it comprises at least 38 amino acid residues, including residues 13-30 of the uPAR-binding site of uPA; competes with labeled DFP-uPA for binding to a cell or molecule that has a binding site for uPA, and has an IC₅₀ value of about 10 nM or less; and is not a fusion protein wherein the uPA peptide is fused to another non-uPA protein or peptide. Preferred molecules are uPA, scuPA, tcuPA, an N-terminal fragment of uPA, residues 1-135, an N-terminal fragment of uPA, residues 1-143, an N-terminal fragment of uPA, residues 1-43; or an N-terminal fragment of uPA, residues 4-43. Detectable labels include a radionuclide, a PET-imageable agent, an MRI-imageable agent, a fluorescer, a fluorogen, a chromophore, a chromogen, a phosphorescer, a chemiluminescer or a bioluminescer. The disclosed methods are used to inhibit cell migration, cell invasion, cell proliferation or angiogenesis, or to induce apoptosis, preferably in the treatment of a subject having a disease or condition associated with undesired cell migration, invasion, proliferation or angiogenesis.

**DIAGNOSTIC PROBES AND THERAPEUTICS TARGETING
uPA AND uPAR**

BACKGROUND OF THE INVENTION

Field of the Invention

5 The present invention relates to conjugates of the urokinase plasminogen activator (uPA) and its fragments that bind to the cell surface receptor uPA (uPAR), the peptides and constructs labeled to deliver a diagnostic probe or a therapeutic agent to the surfaces of cells expressing uPAR. The proteins and peptides of the invention are capable of carrying a suitable detectable or imageable label that will allow qualitative
10 detection and also quantitation of uPAR levels *in vitro* and *in vivo*. Such labeled peptide compositions are therefore useful as diagnostic, prognostic and imaging tools in all diseases and conditions where this receptor plays a pathological or otherwise undesirable role. cDNA probes that detect uPA and uPAR-expressing cells may also be conjugated with similar diagnostic labels and used in *in situ* hybridization assays. Furthermore, by
15 targeting therapeutic agents that "label" the peptide to uPAR, it is possible to achieve a number of biological effects that include cell death, the inhibition of cell movement and migration and the inhibition of angiogenesis.

Description of the Background Art

20 Urokinase-type plasminogen activator (uPA) has been identified as the initiator of several cascades related to tumor progression, invasion and angiogenesis. The uPA system is strongly linked to pathological processes, such as cell invasion and metastasis in cancer (Danø *et al.*, *Adv. Cancer Res.*, 44:139-266 (1985)).

 Cells produce uPA in an inactive form as a 411 amino acid protein, pro-urokinase (pro-uPA) or single-chain uPA (scuPA), which then binds to its receptor,
25 uPAR. This binding event is a prerequisite for the efficient activation of scuPA to two-chain uPA (tcuPA) in a cellular milieu (Ellis *et al.*, *J. Biol. Chem.*, 264:2185-88 (1989)). Pro-uPA is activated by a single proteolytic cleavage between amino acid 158 (Lys) and 159 (Ile) to activate the proenzyme. Cleavage results in the formation of the two-chain active uPA (tcuPA). Cleavage of pro-uPA at the activation site in fact results in a
30 conformational change and in the gain of plasminogen activator activity both with natural and synthetic substrates.

 uPA is a three-domain protein comprising (1) an N-terminal "growth factor domain" (GFD), (2) a kringle domain, and (3) a C-terminal serine protease domain. uPAR, the receptor for pro-uPA, is also a multi-domain protein anchored by a glycosyl-

phosphatidylinositol anchor to the outer leaf of the cell membrane (Behrendt *et al.*, *Biol. Chem. Hoppe-Seyler*, 376:269-279 (1995)).

The amino acid sequence of the N-terminus of human pro-uPA (residues 1-44 of SEQ ID NO:1) is

```

5      Ser Asn Glu Leu His Gln Val Pro Ser Asn Cys Asp Cys Leu Asn Gly
      1                                10
      Gly Thr Cys Val Ser Asn Lys Tyr Phe Ser Asn Ile His Trp Cys Asn
      20                                30
10     Cys Pro Lys Lys Phe Gly Gly Gln His Cys Glu Ile
      40

```

The structure of pro-uPA, SEQ ID NO:1, is shown in Figure 1.

Therapeutic use of uPA, pro-uPA, tissue plasminogen activator (tPA) or streptokinase (for thromboembolism) requires very high dosage due in part to their very rapid clearance. Possible reasons for the short half-life include binding to specific circulating inhibitors, binding to receptors, internalization and degradation of inhibitor-bound and/or receptor-bound PA.

uPAR is not normally expressed at detectable levels on quiescent cells and must therefore be upregulated before it can initiate the activities of the uPA system. uPAR expression is stimulated *in vitro* by differentiating agents such as phorbol esters (Lund *et al.*, *J. Biol. Chem.* 266:5177-5181 (1991)), by the transformation of epithelial cells, and by various growth factors and cytokines such as VEGF, bFGF, HGF, IL-1, TNF α in endothelial cells and by GM-CSF in macrophages (Mignatti *et al.*, *J. Cell Biol.* 113:1193-1201 (1991); Mandriota *et al.*, *J. Biol. Chem.* 270:9709-9716; Yoshida *et al.*, *Inflammation* 20:319-326 (1996)). This up-regulation has the functional consequence of increasing cell motility, invasion, and adhesion (Mandriota *et al.*, *supra*). More importantly, uPAR appears to be up-regulated *in vivo* in most human carcinomas examined to date, specifically, in the tumor cells themselves, in tumor-associated endothelial cells undergoing angiogenesis and in macrophages (Pyke *et al.*, *Cancer Res.* 53:1911-15 (1993) which may participate in the induction of tumor angiogenesis (Lewis *et al.*, *J. Leukoc. Biol.* 57:747-751 (1995)). uPAR expression in cancer patients is present in advanced disease and has been correlated with a poor prognosis in numerous human carcinomas (Hofmann *et al.*, *Cancer* 78:487-92 (1996); Heiss *et al.*, *Nature Med.* 1:1035-39 (1995). Moreover, uPAR is not expressed uniformly throughout a tumor but tends to be associated with the invasive margin and is considered to represent a phenotypic marker of metastasis in human gastric cancer. The fact that uPAR expression is up-regulated only in pathological states involving extracellular

matrix remodeling and cell motility such as cancer makes it an attractive marker for diagnosis as well as a selective target for therapy.

Earlier studies with peptide fragments within the GFD of uPA had shown that residues 20-30 conferred the specificity of binding, but that residues 13-19 were also
5 needed if residues 20-30 were to attain the proper binding conformation. Specifically, the peptide [Ala¹⁹]uPA(12-32), which contains two cysteines (the third cysteine being replaced by Ala to avoid undesired disulfide bond formations) in its open chain form prevented uPA binding to uPAR with an IC₅₀ of 100 nM. In its oxidized cyclic form with an intrachain disulfide bond between Cys¹³ and Cys³¹, the peptide prevented uPA
10 binding with an IC₅₀ of 40 nM. The authors proposed that residues 13-19 might act indirectly to provide a scaffold that would help residues 20-30 attain the correct binding conformation (Appella *et al.*, *J. Biol. Chem.*, 262:4437-4440 (1987).

A related commonly-assigned patent (U.S. 5,942,492) and application (USSN 09/1,816), both of which are incorporated herein by reference in their entirety, show that
15 novel cyclic molecules derived from the uPA peptide fragment 20-30 (in which residue 20 is covalently bonded to residue 30) bind to uPAR with IC₅₀ values in the 10-100 nM range.

Citation of the above documents is not intended as an admission that any of the foregoing is pertinent prior art. All statements as to the date or representation as to the
20 contents of these documents is based on the information available to the applicant and does not constitute any admission as to the correctness of the dates or contents of these documents.

SUMMARY OF THE INVENTION

The present inventors have discovered that relatively large uPA peptides that are
25 cleared rapidly from the body are useful as diagnostic agents (as well as in therapy). These include, but are not limited to, high molecular weight uPA (residues 1-411); single-chain uPA (scuPA) described above; tcuPA, DFP-uPA (tcuPA inactivated with the suicide inhibitor diisopropyl fluorophosphate), the N-terminal fragment often designated ATF, which is a peptide of 135 or 143 amino acids having the sequence of 1-135 or 1-143 of
30 native uPA; the GFD which corresponds to residues 4-43.

The preferred uPAR-targeting protein or peptide has the following characteristics:

- (a) diagnostically or therapeutically labeled (which terms are defined below);
- (b) comprises at least 38 amino acid residues;
- (c) includes residues 13-30 of the uPAR-binding site of uPA.

- (d) competes with labeled DFP-uPA (preferably [¹²⁵I] labeled) for binding to a cell or molecule that has a binding site for uPA, and has an IC₅₀ value of about 10 nM or less -- that is, the protein or peptide causes 50% inhibition of the binding of the labeled ligand at concentrations of 10 nM or lower).
- 5 (e) is not a fusion protein wherein the uPA peptide is fused to another (non-uPA) protein or peptide.

Preferred detectable labels include a radionuclides, PET-imageable agents, MRI-imageable agents, fluorescers, fluorogens, a chromophore, a chromogen, a phosphorescer, a chemiluminescer or a bioluminescer. Such a label permits detection or
10 quantitation of the uPAR level in a tissue sample and can be used, therefore, as a diagnostic and a prognostic tool in all diseases where enhanced expression of the receptor plays a pathological or otherwise undesirable role, including those described herein.

A most preferred radionuclide is selected from the group consisting of ³H, ¹⁴C,
15 ³⁵S, ⁹⁹Tc, ¹²³I, ¹²⁵I, ¹³¹I, ¹¹¹In, ⁹⁷Ru, ⁶⁷Ga, ⁶⁸Ga, ⁷²As, ⁸⁹Zr and ²⁰¹Tl. In the diagnostic composition, the fluorescer or fluorogen is preferably fluorescein, rhodamine, dansyl, phycoerythrin, phycocyanin, allophycocyanin, *o*-phthaldehyde, fluorescamine, a fluorescein derivative, Oregon Green, Rhodamine Green, Rhodol Green or Texas Red.

A protein or peptide probe of this invention, *e.g.*, labeled tcuPA, has the
20 advantage that it may be internalized by the cells to which it binds, *e.g.*, tumor cells. This property is useful for imaging techniques in which it is especially important to reduce background signal (the circulating or non-specifically bound probe molecules) relative to specifically bound probe. This uptake permits clearance of circulating probe so that the ratio of labeled probe inside tumor cells to the probe elsewhere in the body
25 increases. Shorter peptides, such as those described in the Background section, would not be expected to behave in this way. Thus, the present invention provides agents with increased "contrast" for detecting tumors or other sites of cells that express uPAR.

The present invention provides the first specifically-targeted contrast agent for magnetic resonance imaging (MRI). The specificity is a result of the enhanced
30 expression of uPAR in the target tissue, generally tumor tissue. Some of the peptides that bind uPAR and, as a result undergo activation, may be quenched by the naturally occurring plasminogen activator inhibitor-1 (PAI-1) and are taken up by a uPAR-dependent mechanism as uPA peptide-PAI-1 complexes.

Preferably, a diagnostic label is bound to the protein or peptide through one or
35 more diethylenetriaminepentaacetic acid (DTPA) residues that are coupled to the protein

or peptide. In a preferred embodiment, where labeling is targeted, the label is bound through one DTPA residue. A most preferred diagnostic peptides is scuPA or ATF 1-135 coupled to one (or more) DTPA residues, to which are bound gadolinium atoms. Preferred diagnostic methods comprise MRI using these labeled peptides.

5 In another embodiment, the protein or peptide carries a suitable therapeutic "label" also referred to herein as a "therapeutic moiety." A therapeutic moiety is an atom, a molecule, a compound or any chemical component added to the peptide that renders it active in treating a disease or condition associated with enhanced expression of uPAR. The peptides of the present invention are prepared by conventional means, either
 10 recombinant or by proteolysis of uPA as described in Mazar *et al.*, (*Fibrinolysis 6 (suppl. 1)*: 49-55), hereby incorporated by reference in its entirety. The therapeutically active moiety may be bound directly or indirectly to the peptide. The therapeutic composition may be one in which the therapeutic moiety is bound to the active site of the uPA protein or peptide; such compositions are made from compounds described
 15 below. The therapeutically labeled protein or peptide is administered as pharmaceutical composition which comprises a pharmaceutically acceptable carrier or excipient, and is preferably in a form suitable for injection.

In the above pharmaceutical composition, the therapeutically active moiety is preferably a radionuclide. Examples of radionuclides include ^{47}Sc , ^{67}Cu , ^{90}Y , ^{109}Pd , ^{125}I ,
 20 ^{131}I , ^{186}Re , ^{188}Re , ^{199}Au , ^{211}At , ^{212}Pb and ^{217}Bi .

The present invention also provides a uPA active site-targeting compound that covalently modifies the active site of tPA or a fragment or subunit thereof, which fragment or subunit retains (i) the uPA enzymatic endosite and (ii) a uPAR-binding epitope. This compound may include (a) a detectable label; (b) a therapeutic moiety; or
 25 (c) a chelator that is optionally bound to a detectable label or a therapeutic moiety. The compound localizes said chelator, detectable label or therapeutic moiety to the uPA active site.

The above compound may be an affinity label or a uPA-activated irreversible inhibitor. Preferred affinity labels are alkylating groups, such as chloromethylketone
 30 (CMK).

The above compound preferably has a structure characterized by one of the following four general formulas:

(Label)-(Xaa)₂₋₆-(Lys,Arg)-(alkylating group);
 (Therapeutic moiety)-(Xaa)₂₋₆-(Lys,Arg)-(alkylating group);
 35 (Chelator_(empty))-(Xaa)₂₋₆-(Lys,Arg)-(alkylating group); or
 (Label-Chelator)-(Xaa)₂₋₆-(Lys,Arg)-(alkylating group).

where Xaa is any amino acid and the "label" is a detectable label. In these formulas and throughout the specification, the expression (Lys,Arg) means a single amino acid that is either Lys or Arg. Preferably (Xaa)₂₋₆ is Glu-Gly, resulting in compounds of the formula:

- 5 (Chelator_(empty))-Glu-Gly-Arg-CMK; or
(Label-Chelator)- Glu-Gly-Arg-CMK.

Also provided is a molecule, comprising uPA, tcuPA or a fragment or subunit thereof, which fragment or subunit retains (i) the uPA enzymatic endosite and (ii) a uPAR-binding epitope, and which uPA, tcuPA, fragment or subunit is bonded covalently
10 to the compound described above. This molecule would have the general formula:

(Chelator_(empty))-(Xaa)₂₋₆-(Lys,Arg)-(alkylating group)-uPA; or
(Label-Chelator)-(Xaa)₂₋₆-(Lys,Arg)-(alkylating group)-uPA.

Another embodiment of a uPA active site-targeting peptide compound is one that binds to the endosite and one or more exosites of tcuPA or of a tcuPA
15 fragment or subunit (which fragment or subunit retains the uPA (i) enzymatic endosite and (ii) a uPAR-binding epitope). The peptide compound, which covalently modifies the endosite, preferably includes either (a) detectable label, (b) a therapeutic moiety, or (c) a chelator that is optionally bound to a detectable label or to a therapeutic moiety. This peptide compound localizes said chelator,
20 detectable label or therapeutic moiety to the uPA active site.

The above peptide compound of claim preferably has a structure defined by one of the following four general formulas

- (Label)-(Peptide Z)-(Xaa)₂₋₆-(Lys,Arg)-(alkylating group);
(Therapeutic Moiety)-(Peptide Z)-(Xaa)₂₋₆-(Lys,Arg)-(alkylating group),
25 (Chelator_(empty))-(Peptide Z)-(Xaa)₂₋₆-(Lys,Arg)-(alkylating group); or
(Label-Chelator)-(Peptide Z)-(Xaa)₂₋₆-(Lys,Arg)-(alkylating group).

The (Label) is a detectable label, Xaa is any amino acid and Peptide Z is any peptide that binds to a surface exosite of uPA.

Also included is a molecule, comprising uPA, tcuPA or a fragment or subunit of
30 tcuPA bonded at the uPA endosite and at one or more exosites to the above peptide compound.

In the above method, the protein or peptide is preferably scuPA or said N-terminal fragment of uPA residues 1-135 and the detectable label is preferably gadolinium.

The invention includes methods of using the above classes of molecules, specifically, in a method for detecting the presence of uPAR (i) on the surface of a cell, (ii) in a tissue, (iii) in an organ or (iv) in a biological sample, which cell, tissue, organ or sample is suspected of expressing uPAR due to a pathological state. The method comprising the steps of (a) contacting the cell, tissue, organ or sample with any of the diagnostic and detectably labeled compositions and molecules described above; and (b) detecting the presence of the label associated with the cell, tissue, organ or sample. In this method, both the contacting and the detecting may be conducted *in vitro*. Alternatively, the contacting is *in vivo* and the detecting is *in vitro*. Preferably, the contacting and the detecting are *in vivo*.

A nonlimiting group of diseases or conditions treatable by therapeutically labeled peptides of this invention include primary growth of solid tumors or leukemias and lymphomas, metastasis, invasion and/or growth of tumor metastases.

The present invention also provides a method for inhibiting cell migration, cell invasion, cell proliferation or angiogenesis, or for inducing apoptosis, comprising contacting cells associated with undesired cell migration, invasion, proliferation or angiogenesis with an effective amount of the above therapeutically labeled composition. Preferably, the method is used to inhibit the invasiveness of tumor cells.

Diseases or conditions treatable by the present composition include primary growth of a solid tumor, leukemia or lymphoma; tumor invasion, metastasis or growth of tumor metastases; benign hyperplasia; atherosclerosis; myocardial angiogenesis; post-balloon angioplasty vascular restenosis; neointima formation following vascular trauma; vascular graft restenosis; coronary collateral formation; deep venous thrombosis; ischemic limb angiogenesis; telangiectasia; pyogenic granuloma; corneal disease; rubeosis; neovascular glaucoma; diabetic and other retinopathy; retrolental fibroplasia; diabetic neovascularization; macular degeneration; endometriosis; arthritis; fibrosis associated with a chronic inflammatory condition, traumatic spinal cord injury including ischemia, scarring or fibrosis; lung fibrosis, chemotherapy-induced fibrosis; wound healing with scarring and fibrosis; peptic ulcers; a bone fracture; keloids; or a disorder of vasculogenesis, hematopoiesis, ovulation, menstruation, pregnancy or placentation associated with pathogenic cell invasion or with angiogenesis.

Another preferred use of the present methods is the hybridization of a fluorescently labeled cDNA probe (FISH probe) complementary to uPA or to uPAR mRNA expressed in a tumor and detected *in vitro* in a paraffin embedded or frozen tissue section derived from that tumor.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 is a schematic representation of the pro-uPA molecule (SEQ ID NO:1]. The N-terminal growth factor domain (ATF) of human uPA is residues 1-135.

DESCRIPTION OF THE PREFERRED EMBODIMENTS

5 The present inventors have discovered that relatively large uPA fragments or peptides that are capable of binding to uPAR, a receptor expressed in pathological conditions such as tumors, can be modified to incorporate diagnostic or therapeutic labels without significant alteration in their binding properties

uPA-uPAR interactions may exhibit varying degrees of species specificity.

10 Thus, it is always preferred to use a peptide having the amino acid sequence (or a substitution variant thereof) of the species of the animal being targeted.

In Vitro Testing of Compositions

A. Assay for ligand binding to uPAR on whole cells

The uPAR-targeting peptide compounds of the invention are readily tested for
15 their binding to uPAR, preferably by measuring their ability to inhibit the binding of [¹²⁵I]DFP-uPA to uPAR in a competitive ligand-binding assay. The assay may employ whole cells that express uPAR, for example cell lines such as RKO or HeLa. A preferred assay is conducted as follows. Cells (about 5 x 10⁴/well) are plated in medium (e.g., MEM with Earle's salts/10% FBS + antibiotics) in 24-well plates, then incubated
20 in a humid 5% CO₂ atmosphere until the cells reach 70% confluence. Catalytically inactivated high molecular weight uPA (DFP-uPA) is radioiodinated using Iodo-gen[®] (Pierce) to a specific activity of about 250,000 cpm/mg. The cell-containing plates are then chilled on ice and the cells are washed twice (5 minutes each) with cold PBS/ 0.05% Tween-80. Test compounds are serially diluted in cold PBS/ 0.1 % BSA/ 0.01%
25 Tween-80 and added to each well to a final volume of 0.3mL 10 minutes prior to the addition of the [¹²⁵I]DFP-uPA. Each well then receives 9500 cpm of [¹²⁵I]DFP-uPA at a final concentration of 0.2 nM). The plates are then incubated at 4°C for 2 hrs, after which time the cells are washed 3x (5 minutes each) with cold PBS/ 0.05% Tween-80. NaOH (1N) is added to each well in 0.5 mL to lyse the cells, and the plate is incubated
30 for 5 minutes at room temperature or until all the cells in each well are lysed as determined by microscopic examination. The contents of each well are then aspirated and the total counts in each well determined using a gamma counter. Each compound is tested in triplicate and the results are expressed as a percentage of the total radioactivity

measured in wells containing [125 I]DFP-uPA alone, which is taken to represent maximum (100 %) binding.

The inhibition of binding of [125 I]DFP-uPA to uPAR is usually dose-related, such that the concentration of the test compound necessary to produce a 50% inhibition of binding (the IC_{50} value), which is expected to fall in the linear part of the curve, is easily determined. In general, the compounds of the invention have IC_{50} values of less than about 10^{-7} M. Preferably, the compounds of the invention have IC_{50} values of less than about 10^{-8} M and, even more preferably, less than about 10^{-9} M.

In Vivo Study of the uPAR-Targeted Peptides

10 A. Xenograft model of subcutaneous (s.c.) tumor growth

Nude mice are inoculated with MDA-MB-231 cells (human breast carcinoma) and Matrigel® (1 x 10^6 cells in 0.2mL) s.c. in the right flank of the animals. The tumors are staged to 100 mm³ and then imaging with a test composition is initiated (50 µg/animal i.p.). Animals are subjected to the appropriate scanning or imaging method
15 when tumors are expected to be at a range of volumes.

B. Xenograft Model of Metastasis

The compounds of this invention can also diagnose the presence of late metastasis using an experimental metastasis model (Crowley, C.W. *et al.*, *Proc. Natl. Acad. Sci. USA* 90 5021-5025 (1993)). Late metastasis involves the steps of attachment
20 and extravasation of tumor cells, local invasion, seeding, proliferation and angiogenesis. Human prostatic carcinoma cells (PC-3) are inoculated into nude mice i.v. via the tail vein and metastases identified after about 14 days, particularly in the lungs but also in regional lymph nodes, femurs and brain. This mimics the organ tropism of naturally occurring metastases in human prostate cancer. A compound of the invention (50
25 µg/mL) is given i.v. on day 14 and the microscopic metastases are imaged using the appropriate detection method. Tissues may also be harvested from these mice after euthanasia followed by the detection of cells expressing uPA and uPAR using the cDNA probes.

For a compound to be useful in accordance with this invention, it should
30 demonstrate an ability to detect the presence of a tumor mass (primary or metastatic) as small as about 5 mm in a mouse model.

Diagnostic and Prognostic Compositions

The uPA peptides can be detectably labeled and used, for example, to detect a peptide binding site or receptor (such as uPAR) on the surface or in the interior of a cell.
35 The fate of the peptide during and after binding can be followed *in vitro* or *in vivo* by

using the appropriate method to detect the label. The labeled peptide may be utilized *in vivo* for diagnosis and prognosis, for example to image occult metastatic foci or for other types of *in situ* evaluations.

A number of U.S. patents, incorporated by reference herein, disclose methods and compositions for complexing metals to larger molecules, including description of useful chelating agents. The metals are preferably detectable metal atoms, including radionuclides, and are complexed to proteins and other molecules. These documents include: US 5,627,286 (Heteroatom-bearing ligands and metal complexes thereof); US 5,618,513 (Method for preparing radiolabeled peptides); US 5,567,408 (YIGSR peptide radiopharmaceutical applications); US 5,443,816 (Peptide-metal ion pharmaceutical preparation and method); US 5,561,220 (Technetium-^{99m} labeled peptides for imaging inflammation).

Suitable detectable labels include radioactive, fluorescent, fluorogenic, chromogenic, or other chemical labels. Useful radiolabels, which are detected simply by gamma counter, scintillation counter or autoradiography include ³H, ¹²⁵I, ¹³¹I, ³⁵S and ¹⁴C. In addition, ¹³¹I is a useful therapeutic isotope (see below).

Common fluorescent labels include fluorescein, rhodamine, dansyl, phycoerythrin, phycocyanin, allophycocyanin, *o*-phthaldehyde and fluorescamine. The fluorophore, such as the dansyl group, must be excited by light of a particular wavelength to fluoresce. See, for example, Haugland, *Handbook of Fluorescent Probes and Research Chemicals*, Sixth Ed., Molecular Probes, Eugene, OR., 1996). Fluorescein, fluorescein derivatives and fluorescein-like molecules such as Oregon GreenTM and its derivatives, Rhodamine GreenTM and Rhodol GreenTM, are coupled to amine groups using the isothiocyanate, succinimidyl ester or dichlorotriazinyl-reactive groups. Similarly, fluorophores may also be coupled to thiols using maleimide, iodoacetamide, and aziridine-reactive groups. The long wavelength rhodamines, which are basically Rhodamine GreenTM derivatives with substituents on the nitrogens, are among the most photostable fluorescent labeling reagents known. Their spectra are not affected by changes in pH between 4 and 10, an important advantage over the fluoresceins for many biological applications. This group includes the tetramethylrhodamines, X-rhodamines and Texas RedTM derivatives. Other preferred fluorophores for derivatizing the peptide according to this invention are those which are excited by ultraviolet light. Examples include cascade blue, coumarin derivatives, naphthalenes (of which dansyl chloride is a member), pyrenes and pyridyloxazole derivatives. Also included as labels are two related inorganic materials that have recently been described: semiconductor

nanocrystals, comprising, for example, cadmium sulfate (Bruchez, M. *et al.*, *Science* 281:2013-2016 (1998), and quantum dots, *e.g.*, zinc-sulfide-capped cadmium selenide (Chan, W.C.W. *et al.*, *Science* 281:2016-2018 (1998)).

In yet another approach, the amino group of a uPA peptide is allowed to react
 5 with reagents that yield fluorescent products, for example, fluorescamine, dialdehydes such as *o*-phthaldialdehyde, naphthalene-2,3-dicarboxylate and anthracene-2,3-dicarboxylate. 7-nitrobenz-2-oxa-1,3-diazole (NBD) derivatives, both chloride and fluoride, are useful to modify amines to yield fluorescent products.

The peptides can also be labeled for detection using fluorescence-emitting metals
 10 such as ^{152}Eu , or others of the lanthanide series. These metals can be attached to the peptide using such metal chelating groups as diethylenetriaminepentaacetic acid (DTPA) or ethylenediaminetetraacetic acid (EDTA). DTPA, for example, is available as the anhydride, which can readily modify the NH_2 -containing uPAR-binding peptides of this invention.

For *in vivo* diagnosis, radionuclides may be bound to the peptide either directly
 15 or indirectly using a chelating agent such as DTPA and EDTA which is chemically conjugated, coupled or bound (which terms are used interchangeably) to the peptide. The chemistry of chelation is well known in the art, and a varying range of molar ratios of DTPA:peptide may be used to produce the DTPA-coupled peptide, and thereby, the
 20 diagnostically labeled peptide. The key limiting factor on the chemistry of coupling is that the uPA fragment must maintain its ability to bind uPAR after labeling. The term "chelating agent" and "chelator" and "chelating group" are meant to be synonymous and to be distinct from "chelate" which is the formed complex comprising the chelator and the metal atom or ion.

Any radionuclide having diagnostic or therapeutic value can be used as the
 25 radiolabel in the present invention. In a preferred embodiment, the radionuclide is a γ -emitting or beta-emitting radionuclide, for example, one selected from the lanthanide or actinide series of the elements. Positron-emitting radionuclides, *e.g.* ^{68}Ga or ^{64}Cu , may also be used. Suitable gamma-emitting radionuclides include those which are useful in
 30 diagnostic imaging applications. The gamma-emitting radionuclides preferably have a half-life of from 1 hour to 40 days, preferably from 12 hours to 3 days. Examples of suitable gamma-emitting radionuclides include ^{67}Ga , ^{111}In , $^{99\text{m}}\text{Tc}$, ^{169}Yb and ^{186}Re . Most preferably, the radionuclide is $^{99\text{m}}\text{Tc}$.

Examples of preferred radionuclides (ordered by atomic number) are ^{67}Cu , ^{67}Ga ,
 35 ^{68}Ga , ^{72}As , ^{89}Zr , ^{90}Y , ^{97}Ru , ^{99}Tc , ^{111}In , ^{123}I , ^{125}I , ^{131}I , ^{169}Yb , ^{186}Re , and ^{201}Tl . Most preferred

are gamma-emitting radionuclides such as ^{67}Ga , ^{111}In , $^{99\text{m}}\text{Tc}$, ^{169}Yb and ^{186}Re , most preferably $^{99\text{m}}\text{Tc}$. Though limited work have been done with positron-emitting radiometals as labels, certain proteins, such as transferrin and human serum albumin, have been labeled with ^{68}Ga

5 A number of metals (not radioisotopes) useful for MRI include gadolinium, manganese, copper, iron, gold and europium. Gadolinium is most preferred. Generally, the amount of labeled peptide needed for detectability in diagnostic use will vary depending on considerations such as age, condition, sex, and extent of disease in the patient, contraindications, if any, and other variables, and is to be adjusted by the
10 individual physician or diagnostician. Dosage can vary from 0.01 mg/kg to 100 mg/kg.

 The peptides can also be made detectable by coupling them to a phosphorescent or a chemiluminescent compound. The presence of the chemiluminescent-tagged peptide is then determined by detecting the presence of luminescence that arises during the course of a chemical reaction. Examples of particularly useful chemiluminescers are
15 luminol, isoluminol, theromatic acridinium ester, imidazole, acridinium salt and oxalate ester. Likewise, a bioluminescent compound may be used to label the peptides. Bioluminescence is a type of chemiluminescence found in biological systems in which a catalytic protein increases the efficiency of the chemiluminescent reaction. The presence of a bioluminescent protein is determined by detecting the presence of luminescence.
20 Important bioluminescent compounds for purposes of labeling are luciferin, luciferase and aequorin.

 In yet another embodiment, colorimetric detection is used, based on chromogenic compounds which have, or result in, chromophores with high extinction coefficients.

In situ detection of the labeled peptide may be accomplished by removing a
25 histological specimen from a subject and examining it by microscopy under appropriate conditions to detect the label. Those of ordinary skill will readily perceive that any of a wide variety of histological methods (such as staining procedures) can be modified in order to achieve such *in situ* detection.

 The term "diagnostically labeled" means that the peptide has attached to it a
30 diagnostically detectable label. There are many different labels and methods of labeling known to those of ordinary skill in the art. Examples of the types of labels which can be used in the present invention include radioactive isotopes, paramagnetic isotopes, and compounds which can be imaged by positron emission tomography (PET) and by MRI. Those of ordinary skill in the art will know of other suitable labels for binding to the

peptides used in the invention, or will be able to ascertain such, by routine experimentation.

For diagnostic *in vivo* radioimaging, the type of detection instrument available is a major factor in selecting a radionuclide. The radionuclide chosen must have a type of decay which is detectable by a particular instrument. In general, any conventional method for visualizing diagnostic imaging can be utilized in accordance with this invention. Another factor in selecting a radionuclide for *in vivo* diagnosis is that its half-life be long enough so that the label is still detectable at the time of maximum uptake by the target tissue, but short enough so that deleterious irradiation of the host is minimized. In one preferred embodiment, a radionuclide used for *in vivo* imaging does not emit particles, but produces a large number of photons in a 140-200 keV range, which may be readily detected by conventional gamma cameras.

In vivo imaging may be used to detect occult metastases which are not observable by other methods. The expression of uPAR correlates with progression of diseases in cancer patients such that patients with late stage cancer have higher levels of uPAR in both their primary tumors and metastases. uPAR-targeted imaging could be used to stage tumors non-invasively or to detect other diseases which are associated with the presence of increased levels of uPAR (for example, restenosis that occurs following angioplasty).

The compositions of the present invention may be used in diagnostic, prognostic or research procedures in conjunction with any appropriate cell, tissue, organ or biological sample of the desired animal species. By the term "biological sample" is intended any fluid or other material derived from the body of a normal or diseased subject, such as blood, serum, plasma, lymph, urine, saliva, tears, cerebrospinal fluid, milk, amniotic fluid, bile, ascites fluid, pus and the like. Also included within the meaning of this term is a organ or tissue extract and a culture fluid in which any cells or tissue preparation from the subject has been incubated.

The labeled peptide compounds of the invention, as well as diagnostically acceptable salts thereof, may be incorporated into convenient dosage forms

Preferably, the compounds of the invention are administered systemically, *e.g.*, by injection. When used, injection may be by any known route, preferably intravenous, subcutaneous, intramuscular, intracranial or intraperitoneal. Injectables can be prepared in conventional forms, either as solutions or suspensions, solid forms suitable for solution or suspension in liquid prior to injection, or as emulsions.

Solid carriers include starch, lactose, calcium sulfate dihydrate, terra alba, sucrose, talc, gelatin, agar, pectin, acacia, magnesium stearate and stearic acid. Liquid carriers include syrup, peanut oil, olive oil, saline, water, dextrose, glycerol and the like. Similarly, the carrier or diluent may include any prolonged release material, such as glyceryl monostearate or glyceryl distearate, alone or with a wax. When a liquid carrier is used, the preparation may be in the form of a syrup, elixir, emulsion, soft gelatin capsule, sterile injectable liquid (e.g., a solution), such as an ampoule, or an aqueous or nonaqueous liquid suspension. A summary of such pharmaceutical compositions may be found, for example, in *Remington's Pharmaceutical Sciences*, Mack Publishing Company, Easton Pennsylvania (Gennaro 18th ed. 1990).

The present invention may be used in the diagnosis of any of a number of animal genera and species, and are equally applicable in the practice of human or veterinary medicine. Thus, the compositions can be used with domestic and commercial animals, including birds and more preferably mammals, as well as humans.

The peptides in a form suitable for topic application are sprayable aerosol preparations wherein the compound, preferably in combination with a solid or liquid inert carrier material, is packaged in a squeeze bottle or in admixture with a pressurized volatile, normally gaseous propellant. The aerosol preparations can contain solvents, buffers, surfactants, perfumes, and/or antioxidants in addition to the compounds of the invention.

An alternative diagnostic approach utilizes cDNA probes that are complementary to and thereby detect uPA and uPAR-expressing cells by *in situ* hybridization with mRNA in these cells. This cDNA is preferably fluorescently labeled. This method and the preparations useful therein are described below.

Active Site-Specific Modification of uPA and other Serine Proteases

The present invention is directed to an active site-specific reagent that covalently modifies the active site ("endsite") of uPA or any serine protease and localizes a detectable label, preferably an imageable label, or a therapeutic moiety such as a radioisotope, toxin or drug conjugate, to the endsite. To be useful in this embodiment, the uPA must be tcuPA or a similar molecule that retains the enzymatic endsite of uPA and its uPAR-binding epitope. Such a reagent is preferably an active site-directed affinity label or an enzyme-activated irreversible inhibitor of uPA activity ("suicide substrate"). In one embodiment, the reagent is attached directly (*i.e.*, without the mediation of a chelating agent) to a detectable label or therapeutic moiety. In another embodiment, the reagent includes as part of its structure a chelating agent, as described above, that is either loaded with the label (or therapeutic moiety) or to which the label

(or therapeutic moiety) may be added after the chelator has been delivered to the enzyme's endosite. The endosite-specific reagent is preferably a modified peptide, but may also be a non-peptidic molecule, as described below.

The molecules and approaches described herein are useful for other serine proteases, which utilize a similar catalytic mechanism as uPA based on the canonical catalytic triad of Ser, His and Asp common to all serine proteases. Thus, any active site-directed affinity label or suicide substrate of any serine protease can be used to localize to the endosite of that protease a chelating agent or a chelate that includes a detectable (preferably imageable) label or therapeutic moiety.

A typical example of such a reagent is an endosite-specific affinity label comprising a tripeptide and any alkylating moiety. Upon being localized to the active site of a serine protease, the reagent reacts with the active site histidine. A preferred alkylating moiety is a halomethylketone, more preferably chloromethylketone ("CMK"). A preferred endosite-specific tripeptide for uPA is Glu-Gly-Arg. Thus a preferred reagent is Glu-Gly-Arg-CMK to which is bonded a label, a therapeutic moiety or a chelating agent optionally loaded with a label or therapeutic moiety. This reagent molecule targets the His residue within the serine protease catalytic triad of uPA and covalently bonds to this His residue. Other peptides that may be longer than a tripeptide are also useful – they may have even higher affinities or be more reactive than Glu-Gly-Arg for the protease endosite and could also access exosites, thereby resulting in increased binding affinity and specificity. Bock reported the incorporation of spectroscopic probes into active sites of certain serine proteases by inactivating the enzyme with a thioester derivative of a peptide CHK. The thiol group that is generated can serve as a unique suite for subsequent labeling with thiol-reactive probes, exemplified by 5-(iodoacetamido)fluorescein (Bock, P.E., *Biochemistry* 27:6633-6639 (1999), incorporated by reference in its entirety). Thus, the present approach may be extended along these lines for incorporating the various types of detectable labels or therapeutic moieties not contemplated by Bock for the uses set forth herein.

It is within the skill of the art to test other peptides or similar molecules for binding to the uPA endosite using conventional means, and thereby determine their suitability for use in accordance with this invention. All that is required is the peptide portion of the reagent have a basic residue (Lys or Arg) at the "S1" site. Thus for example, the peptide part of the molecule may have the following structure (where Xaa symbolizes any amino acid residue): (Xaa)₂₋₆-(Lys,Arg), wherein the (Lys,Arg) is intended to be either Lys or Arg. A preferred reagent would be

(Xaa)₂₋₆-(Lys,Arg)-CMK. Furthermore, non-peptide affinity labels or suicide substrates that inactivate a serine protease active site could also be useful in localizing to the enzyme active site (1) a detectable label or therapeutic moiety; or (2) a chelate that contains, or a chelator that has the potential to contain, a detectable label or therapeutic moiety. These non-peptide reagents are not limited to those that modify the catalytic triad His residue, as compounds are known which can alkylate the active site Ser (*e.g.*, diisopropyl fluorophosphate, DFP; and phenylmethylsulfonyl fluoride, PMSF). Moreover, Porter, N.A *et al.*, *J. Am Chem Soc.* 121:7716-7717 (1999), incorporated by reference in its entirety, describes the use of photolabile moieties to modify proteases with a photoremovable "cage." Of particular utility herein is the description of 4-aminocinnamate esters to label the endosite of proteases. The present invention includes such types of modifying groups as well.

A reagent containing a chelating group must be capable of modifying a thiol or amine group. In the case of an amine-reactive reagent, because the N-terminal amine has a pK_a of about 7.5 whereas the epsilon amino group of Lys has a pK_a of about 9.5, these two types of amino group will have quite different reactivities within the pH range 7.5 - 9.5. Thus, by varying the pH of the reaction conditions, a reagent delivering the chelating group can be introduced at the N-terminus of a peptidic suicide substrate in a selective manner. The chelating group for an imageable label may be part of the reagent, for example, (Chelator)-(Xaa)₂₋₆-(Lys,Arg)-CMK. It is also possible to introduce a free thiol at the N-terminus of a peptidic suicide inhibitor by introducing a Cys or HomoCys residue as one of the Xaa amino acids or by chemically modifying the N-terminus of the peptide suicide substrate. The chelator may then be part of the reagent, as described above, or it may be introduced into the (Xaa)₂₋₆-(Lys,Arg)-CMK moiety after this peptide reagent has bound to the uPA active site. A more generic characterization of this preferred type of endosite-specific reagent is

(Chelator)-(Xaa)₂₋₆-(Lys,Arg)-(alkylating group).

Once a particular endosite-specific reagent has bound to uPA, a new molecular entity is created which is also intended to be within the scope of this invention. This molecular entity is uPA (or other serine protease), tcuPA, or a fragment thereof that retains the endosite and the uPAR-binding epitope, to which is covalently bonded one of the above reagents. The reagent may incorporate a detectable label, a therapeutic moiety, an "empty" chelator not carrying an imageable label or therapeutic moiety (but capable of accepting one), or a chelator that does carry an imageable label or therapeutic moiety. Alternatively, the chelator, label, *etc.*, is not necessarily incorporated into the

reagent *a priori*; rather, the reagent would be derivatized subsequently to carry the label or moiety. The structures may be characterized as follows (where "label" means detectable label):

- 5 (Label)-(Xaa)₂₋₆-(Lys,Arg)-(alkylating group)-uPA;
 (Therapeutic moiety)-(Xaa)₂₋₆-(Lys,Arg)-(alkylating group)-uPA;
 (Chelator_(empty))-(Xaa)₂₋₆-(Lys,Arg)-(alkylating group)-uPA; or
 (Label-Chelator)-(Xaa)₂₋₆-(Lys,Arg)-(alkylating group)-uPA.


- Serine protease active sites can be fairly restrictive and may only accommodate a few amino acid residues or substrates of limited size (Nozawa *et al.*, *J. Biochem* (1982) 91:1837-1843). The endosite-specific peptide described above does not have a *per se* limitation on its length (although, as indicated, peptides of 3-7 residues are preferred). This absence of a length limitation is because endosite interactions are but one possible way of localizing an alkylating moiety such as CMK to the uPA active site. Thus, another embodiment is a peptide that binds to the uPA endosite and one or more
 15 exosites. An example is an endosite-binding tripeptide which further comprises an additional peptide "tail" that binds to a surface exosite of the protein. A well-known example of such a construct is hirulog, a thrombin inhibitor. Hirulog combines the active site specificity of a thrombin-specific tripeptide and includes more peptide chain that binds to the surface of thrombin, yielding a high-affinity inhibitor of thrombin
 20 activity (Skordalakes *et al. Biochemistry* 13:14420-7).

- Using similar notation to that above, a range of preferred embodiments include:
 (Label)-(Peptide Z)-(Xaa)₂₋₆-(Lys,Arg)-(alkylating group);
 (Therapeutic Moiety)-(Peptide Z)-(Xaa)₂₋₆-(Lys,Arg)-(alkylating group),
 (Chelator_(empty))-(Peptide Z)-(Xaa)₂₋₆-(Lys,Arg)-(alkylating group); and
 25 (Label-Chelator)-(Peptide Z)-(Xaa)₂₋₆-(Lys,Arg)-(alkylating group)
 wherein Peptide Z is any peptide that binds to a surface exosite of uPA.

This would give rise to at least four novel types of molecules shown below, the latter two of which are most preferred:

- In yet another embodiment, a longer peptide (of between about 4 and 30
 30 residues) that mimics the recognition region of PAI-1 is also useful as a site-specific reagent to produce a uPAR-directed imaging protein/peptide as described above.

As has been discussed earlier, other useful reagents modify uPA randomly (not necessarily being localized to the active site) to create uPAR-imaging agents.



In addition to the peptidic reagents described above, a general class of non-
 20 peptidic agents may be used to localize a chelator (and thereby, a detectable label) to the
 active site of uPA or of any serine protease. Such structures include, DFP and PMSF,
etc. A member of this group of compounds should have the following properties:

- (1) it selectively modifies only endosite residues (either the Ser or the His)
- (2) it localizes a chelator or chelate only to the endosite of uPA (or other serine
 25 protease)

These reagents are selective for the active site by virtue of the fact that they are enzyme
 activated irreversible inhibitors and will only form covalent adducts when catalyzed by
 the target enzyme.

Diagnostic Methods

30 The methods of this invention may be used to image tumor foci in a subject. A
 vertebrate subject, preferably a mammal, more preferably a human, is administered an

amount of the compound effective to image a tumor focus. The compound or pharmaceutically acceptable salt thereof is preferably administered in the form of a diagnostic composition as described above.

Useful doses of the compounds are defined as effective amount of the peptide for the particular diagnostic measurement. Thus, an effective amount means an amount sufficient to be detected using the appropriate detection system *e.g.*, MRI detector, gamma camera, *etc.* The minimum detectable amount will depend on the ratio of labeled peptide specifically bound to a tumor (signal) to the amount of labeled peptide either bound non-specifically or found free in plasma or in extracellular fluid. The advantage of the peptides of this invention is that they are cleared rapidly from the plasma and do not accumulate in most tissues, in contrast to antibody-based contrast agents. Thus, the signal-to-noise ratio is increased and only the specifically tumor-bound peptide is detected or quantitated. This allows for accurate localization of the signal and the detection of small tumor lesions which can only be detected in the absence of high background signals.

The amount of the diagnostic composition to be administered depends on the precise peptide selected, the disease or condition, the route of administration, and the judgment of the skilled imaging professional.

A preferred dose for imaging by MRI, SPECT, *etc.*, is an amount of up to about 100 milligrams of labeled peptide compound per kilogram of body weight. A typical single dosage of the labeled peptide is between about 1 ng and about 100 mg/kg body weight.

An alternative diagnostic approach utilizes cDNA probes that are complementary to and thereby detect uPA- and uPAR-expressing cells by *in situ* hybridization with mRNA in these cells. The present invention provides methods for localizing uPAR or uPA mRNA in cells using fluorescent *in situ* hybridization (FISH) with labeled cDNA probes that correspond to part or all of the uPA (or uPAR) coding sequence. The basic principle of FISH is that DNA or RNA in the prepared specimens are hybridized with the probe nucleic acid of this invention that is labeled non-isotopically with, for example, a fluorescent dye, biotin or digoxigenin. The hybridized signals are then detected by fluorimetric or by enzymatic methods, for example, by using a fluorescence or light microscope. The detected signal and image can be recorded on light sensitive film.

An advantage of using a fluorescent probe is that the hybridized image can be readily analyzed using a powerful confocal microscope or an appropriate image analysis

system with a charge-coupled device (CCD) camera. As compared with radioactive methods, FISH offers increased sensitivity. In addition to offering positional information, FISH allows better observation of cell or tissue morphology. Because of the nonradioactive approach, FISH has become widely used for localization of specific DNA or mRNA in a specific cell or tissue type.

The *in situ* hybridization methods and the preparations useful herein are described in Wu, W. *et al.*, eds., *Methods in Gene Biotechnology*, CRC Press, 1997, chapter 13, pages 279-289. This book is incorporated by reference in its entirety, as are the references cited therein. A number of U.S. patents and scientific articles that describe various *in situ* hybridization techniques and applications, also incorporated by reference, are: 5,912,165; 5,906,919; 5,885,531; 5,880,473; 5,871,932; 5,856,097; 5,837,443; 5,817,462; 5,784,162; 5,783,387; 5,750,340; 5,759,781; 5,707,797; 5,677,130; 5,665,540; 5,571,673; 5,565,322; 5,545,524; 5,538,869; and 5,501,954, 5,225,326, 4,888,278. Other related references include Jowett, T., *Methods Cell Biol.* 59:63-85 (1999) Pinkel *et al.*, *Cold Spring Harbor Symp. Quant. Biol.* 51:151-157 (1986); Pinkel, D. *et al.*, *Proc. Natl. Acad. Sci. (USA)* 83:2934-2938 (1986); Gibson *et al.*, *Nucl. Acids Res.* 15:6455-6467 (1987); Urdea *et al.*, *Nucl. Acids Res.* 16:4937-4956 (1988); Cook *et al.*, *Nucl. Acids Res.* 16:4077-4095 (1988); Telser *et al.*, *J. Am. Chem. Soc.* 111:6966-6976 (1989); Allen *et al.*, *Biochemistry* 28:4601-4607 (1989); Nederlof, P.M. *et al.*, *Cytometry* 10:20-27 (1989); Nederlof, P.M. *et al.*, *Cytometry* 11:126-131 (1990); Seibl, R., *et al.*, *Biol. Chem. Hoppe-Seyler* 371:939-951 (Oct. 1990); Wiegant, J. *et al.*, *Nucl. Acids Res.* 19:3237-3241 (1991); McNeil JA *et al.*, *Genet Anal Tech Appl* 8:41-58 (1991); Komminoth *et al.*, *Diagnostic Molecular Biology* 1:85-87 (1992); Dauwerse, J.G. *et al.*, *Hum. Mol. Genet.* 1:593-598 (1992); Ried, T. *et al.*, *Proc. Natl. Acad. Sci. (USA)* 89:1388-1392 (1992); Wiegant, J. *et al.*, *Cytogenet. Cell Genet.* 63:73-76 (1993); Glaser, V., *Genetic. Eng. News.* 16:1, 26 (1996); Speicher, M.R., *Nature Genet.* 12:368-375 (1996)

Pharmaceutical and Therapeutic Compositions and Their Administration

The compounds that may be employed in the pharmaceutical compositions of the invention include all of those compounds described above, as well as the pharmaceutically acceptable salts of these compounds. As stated above, the compounds of the invention possess the ability to inhibit invasiveness or angiogenesis, properties that are exploited in the treatment of cancer, in particular metastatic cancer. A composition of this invention may be active *per se*, or may act as a "pro-drug" that is converted *in vivo* to the active form.

The compounds of the invention, as well as the pharmaceutically acceptable salts thereof, may be incorporated into convenient dosage forms, such as capsules, impregnated wafers, tablets or injectable preparations. Solid or liquid pharmaceutically acceptable carriers may be employed.

5 Preferably, the compounds of the invention are administered systemically, *e.g.*, by injection. When used, injection may be by any known route, preferably intravenous, subcutaneous, intramuscular, intracranial or intraperitoneal. Injectables can be prepared in conventional forms, either as solutions or suspensions, solid forms suitable for solution or suspension in liquid prior to injection, or as emulsions.

10 Solid carriers include starch, lactose, calcium sulfate dihydrate, terra alba, sucrose, talc, gelatin, agar, pectin, acacia, magnesium stearate and stearic acid. Liquid carriers include syrup, peanut oil, olive oil, saline, water, dextrose, glycerol and the like. Similarly, the carrier or diluent may include any prolonged release material, such as glyceryl monostearate or glyceryl distearate, alone or with a wax. When a liquid carrier
15 is used, the preparation may be in the form of a syrup, elixir, emulsion, soft gelatin capsule, sterile injectable liquid (*e.g.*, a solution), such as an ampoule, or an aqueous or nonaqueous liquid suspension. A summary of such pharmaceutical compositions may be found, for example, in *Remington's Pharmaceutical Sciences*, Mack Publishing Company, Easton Pennsylvania (Gennaro 18th ed. 1990).

20 The pharmaceutical preparations are made following conventional techniques of pharmaceutical chemistry involving such steps as mixing, granulating and compressing, when necessary for tablet forms, or mixing, filling and dissolving the ingredients, as appropriate, to give the desired products for oral, parenteral, topical, transdermal, intravaginal, intrapenile, intranasal, intrabronchial, intracranial, intraocular, intraaural
25 and rectal administration. The pharmaceutical compositions may also contain minor amounts of nontoxic auxiliary substances such as wetting or emulsifying agents, pH buffering agents and so forth.

The present invention may be used in the diagnosis or treatment of any of a number of animal genera and species, and are equally applicable in the practice of
30 human or veterinary medicine. The pharmaceutical compositions can therefore be used to treat domestic and commercial animals, including birds and more preferably mammals, as well as humans.

Though the preferred routes of administration are systemic the pharmaceutical composition may be administered topically or transdermally, *e.g.*, as an ointment, cream
35 or gel; orally; rectally; *e.g.*, as a suppository, parenterally, by injection or continuously

by infusion; intravaginally; intrapenilely; intranasally; intrabronchially; intracranially, intraaurally; or intraocularly.

Also suitable for topical application are sprayable aerosol preparations wherein the compound, preferably in combination with a solid or liquid inert carrier material, is packaged in a squeeze bottle or in admixture with a pressurized volatile, normally gaseous propellant. The aerosol preparations can contain solvents, buffers, surfactants, perfumes, and/or antioxidants in addition to the compounds of the invention.

For the preferred topical applications, especially for humans, it is preferred to administer an effective amount of the compound to an affected area, *e.g.*, skin surface, mucous membrane, eyes, *etc.* This amount will generally range from about 0.001 mg to about 1 g per application, depending upon the area to be treated, the severity of the symptoms, and the nature of the topical vehicle employed.

Therapeutic compositions of the invention may comprise, in addition to the labeled peptide, one or more additional anti-tumor agents, such as mitotic inhibitors, *e.g.*, vinblastine; alkylating agents, *e.g.*, cyclophosphamide; folate inhibitors, *e.g.*, methotrexate, piritrexim or trimetrexate; antimetabolites, *e.g.*, 5-fluorouracil and cytosine arabinoside; intercalating antibiotics, *e.g.*, adriamycin and bleomycin; enzymes or enzyme inhibitors, *e.g.*, asparaginase, topoisomerase inhibitors such as etoposide; or biological response modifiers, *e.g.*, interferons or interleukins. In fact, pharmaceutical compositions comprising any known cancer therapeutic in combination with the labeled peptides disclosed herein are within the scope of this invention. The pharmaceutical composition may also comprise one or more other medicaments to treat additional symptoms for which the target patients are at risk, for example, anti-infectives including antibacterial, anti-fungal, anti-parasitic, anti-viral, and anti-coccidial agents.

Therapeutic Compositions

In a preferred embodiment, the peptides describe herein are “therapeutically conjugated” or “therapeutically labeled” (terms which are intended to be interchangeable) and used to deliver a therapeutic agent to the site to which the compounds home and bind, such as sites of tumor metastasis or foci of infection/inflammation, restenosis or fibrosis. The term “therapeutically conjugated” means that the modified peptide is conjugated to another therapeutic agent that is directed either to the underlying cause or to a “component” of tumor invasion, angiogenesis, inflammation or other pathology.

Examples of useful therapeutic radioisotopes (ordered by atomic number) include ⁴⁷Sc, ⁶⁷Cu, ⁹⁰Y, ¹⁰⁹Pd, ¹²⁵I, ¹³¹I, ¹⁸⁶Re, ¹⁸⁸Re, ¹⁹⁹Au, ²¹¹At, ²¹²Pb and ²¹⁷Bi. These atoms can

be conjugated to the peptide directly, indirectly as part of a chelate, or, in the case of iodine, indirectly as part of an iodinated Bolton-Hunter group.

Preferred doses of the radionuclide conjugates are a function of the specific radioactivity to be delivered to the target site which varies with tumor type, tumor location and vascularization, kinetics and biodistribution of the peptide carrier, energy of radioactive emission by the nuclide, *etc.* Those skilled in the art of radiotherapy can readily adjust the dose of the labeled peptide in conjunction with the dose of the particular nuclide to effect the desired therapeutic benefit without undue experimentation.

Another therapeutic approach included here is the use of boron neutron capture therapy (NCT), where a boronated peptide is delivered to a desired target site, such as a tumor, most preferably an intracranial tumor (Barth, R.F., *Cancer Invest.* 14:534-550 (1996); Mishima, Y. (ed.), *Cancer Neutron Capture Therapy*, New York: Plenum Publishing Corp., 1996; Soloway, A.H., *et al.*, (eds), *J. Neuro-Oncol.* 33:1-188 (1997). The stable isotope ^{10}B is irradiated with low energy (<0.025 eV) thermal neutrons, and the resulting nuclear capture yields α particles and ^7Li nuclei which have high linear energy transfer and respective path lengths of about 9 and 5 μm . This method is predicated on ^{10}B accumulation in the tumor with lower levels in blood, endothelial cells and normal tissue (e.g., brain). Such delivery has been accomplished using epidermal growth factor (Yang, W. *et al.*, *Cancer Res* 57:4333-4339 (1997). Because of the selective expression of uPAR in tumors, the peptides of the present invention are excellent delivery vehicles for this therapeutic moiety.

In addition to boron NCT, gadolinium, specifically ^{157}Gd appears to be particularly advantageous for use in NCT with the present peptides. It has recently been reported (Tokumitsu, H. *et al.*, *Chem Pharm Bull* 47:838-842 (1999), incorporated by reference in its entirety) that: ^{157}Gd has the highest thermal neutron capture cross section (255,000 barns) among naturally occurring isotopes, 66 times larger than that of ^{10}B ; Gd neutron capture reaction releases the long range ($>100\mu\text{m}$) prompt γ -rays, internal conversion electrons, X-rays and Auger electrons. Thus, Gd-NCT may increase the chance for photons to hit tumor cells and for electrons to damage these cell locally and intensively. Another advantage is that Gd has long been used as a MRI imaging diagnostic agent. It will be possible to integrate Gd-NCT with MRI diagnosis by using the Gd-loaded dosage forms of the present peptides. A preferred form of Gd for labeling the peptides of this invention for use in Gd-NCT is gadopentetic acid (Gd-DTPA).

Other therapeutic agents which can be coupled to the peptide compounds according to the method of the invention are drugs, prodrugs, enzymes for activating pro-drugs, photosensitizing agents, gene therapeutics, antisense vectors, viral vectors, lectins and other toxins.

5 The therapeutic dosage administered is an amount that is therapeutically effective, as is known to or readily ascertainable by those skilled in the art. The dose is also dependent upon the age, health, and weight of the recipient, kind of concurrent treatment(s), if any, the frequency of treatment, and the nature of the effect desired, such as, for example, anti-inflammatory effects or anti-bacterial effect.

10 Lectins are proteins, commonly derived from plants, that bind to carbohydrates. Among other activities, some lectins are toxic. Some of the most cytotoxic substances known are protein toxins of bacterial and plant origin (Frankel, A.E. *et al.*, *Ann. Rev. Med.* 37:125-142 (1986)). These molecules binding the cell surface and inhibit cellular protein synthesis. The most commonly used plant toxins are ricin and abrin; the most commonly
15 used bacterial toxins are diphtheria toxin and Pseudomonas exotoxin A. In ricin and abrin, the binding and toxic functions are contained in two separate protein subunits, the A and B chains. The ricin B chain binds to the cell surface carbohydrates and promotes the uptake of the A chain into the cell. Once inside the cell, the ricin A chain inhibits protein synthesis by inactivating the 60S subunit of the eukaryotic ribosome Endo, Y. *et al.*, *J.*
20 *Biol. Chem.* 262: 5908-5912 (1987)). Other plant derived toxins, which are single chain ribosomal inhibitory proteins, include pokeweed antiviral protein, wheat germ protein, gelonin, dianthins, momorcharins, trichosanthin, and many others (Strip, F. *et al.*, *FEBS Lett.* 195:1-8 (1986)). Diphtheria toxin and Pseudomonas exotoxin A are also single chain proteins, and their binding and toxicity functions reside in separate domains of the same
25 protein chain with full toxin activity requiring proteolytic cleavage between the two domains. Pseudomonas exotoxin A has the same catalytic activity as diphtheria toxin. Ricin has been used therapeutically by binding its toxic α -chain, to targeting molecules such as antibodies to enable site-specific delivery of the toxic effect. Bacterial toxins have also been used as anti-tumor conjugates. As intended herein, a toxic peptide chain or
30 domain is conjugated to a compound of this invention and delivered in a site-specific manner to a target site where the toxic activity is desired, such as a metastatic focus. Conjugation of toxins to protein such as antibodies or other ligands are known in the art (Olsnes, S. *et al.*, *Immunol. Today* 10:291-295 (1989); Vitetta, E.S. *et al.*, *Ann. Rev. Immunol.* 3:197-212 (1985)).

Cytotoxic drugs that interfere with critical cellular processes including DNA, RNA, and protein synthesis, have been conjugated to antibodies and subsequently used for in vivo therapy. Such drugs, including, but not limited to, daunorubicin, doxorubicin, methotrexate, and mitomycin C are also coupled to the compounds of this invention and used therapeutically in this form.

In another embodiment of the invention, photosensitizers may be coupled to the compounds of the invention for delivery directly to a tumor.

Therapeutic Methods

The methods of this invention may be used to inhibit tumor growth and invasion in a subject or to suppress angiogenesis induced by tumors by inhibiting endothelial cell growth and migration. By inhibiting the growth or invasion of a tumor or angiogenesis, the methods result in inhibition of tumor metastasis. A vertebrate subject, preferably a mammal, more preferably a human, is administered an amount of the compound effective to inhibit tumor growth, invasion or angiogenesis. The compound or pharmaceutically acceptable salt thereof is preferably administered in the form of a pharmaceutical composition as described above.

Doses of the compounds preferably include pharmaceutical dosage units comprising an effective amount of the peptide. By an effective amount is meant an amount sufficient to achieve a steady state concentration *in vivo* which results in a measurable reduction in any relevant parameter of disease and may include growth of primary or metastatic tumor, any accepted index of inflammatory reactivity, or a measurable prolongation of disease-free interval or of survival. For example, a reduction in tumor growth in 20 % of patients is considered efficacious (Frei III, E., *The Cancer Journal* 3:127-136 (1997)). However, an effect of this magnitude is not considered to be a minimal requirement for the dose to be effective in accordance with this invention.

In one embodiment, an effective dose is preferably 10-fold and more preferably 100-fold higher than the 50% effective dose (ED₅₀) of the compound in an *in vivo* assay as described herein.

The amount of active compound to be administered depends on the precise peptide or derivative selected, the disease or condition, the route of administration, the health and weight of the recipient, the existence of other concurrent treatment, if any, the frequency of treatment, the nature of the effect desired, for example, inhibition of tumor metastasis, and the judgment of the skilled practitioner.

A preferred dose for treating a subject, preferably mammalian, more preferably human, with a tumor is an amount of up to about 100 milligrams of active compound per kilogram of body weight. A typical single dosage of the peptide is between about 1 ng and about 100mg/kg body weight. For topical administration, dosages in the range of about 0.01-20% concentration (by weight) of the compound, preferably 1-5%, are suggested. A total daily dosage in the range of about 0.1 milligrams to about 7 grams is preferred for intravenous administration. The foregoing ranges are, however, suggestive, as the number of variables in an individual treatment regime is large, and considerable excursions from these preferred values are expected.

An effective amount or dose of the peptide for inhibiting invasion *in vitro* is in the range of about 1 picogram to about 0.5 nanograms per cell. Effective doses and optimal dose ranges may be determined *in vitro* using the methods described herein.

The foregoing compositions and treatment methods are useful for inhibiting cell migration and invasion or cell proliferation in a subject having any disease or condition associated with undesired cell invasion, proliferation, angiogenesis or metastasis. Such diseases or conditions may include primary growth of solid tumors or leukemias and lymphomas, metastasis, invasion and/or growth of tumor metastases, benign hyperplasias, atherosclerosis, myocardial angiogenesis, angiofibroma, arteriovenous malformations, post-balloon angioplasty vascular restenosis, neointima formation following vascular trauma, vascular graft restenosis, coronary collateral formation, deep venous thrombosis, ischemic limb angiogenesis, telangiectasia, pyogenic granuloma, corneal diseases, rubeosis, neovascular glaucoma, diabetic and other retinopathy, retrolental fibroplasia, diabetic neovascularization, macular degeneration, endometriosis, arthritis, fibrosis associated with chronic inflammatory conditions including psoriasis scleroderma, hemangioma, hemophilic joints, hypertrophic scars, Osler-Weber syndrome, psoriasis, pyrogenic granuloma, retrolental fibroplasia, scleroderma, Von-Hippel-Landau syndrome, trachoma, vascular adhesions, lung fibrosis, chemotherapy-induced fibrosis, wound healing with scarring and fibrosis, peptic ulcers, fractures, keloids, and disorders of vasculogenesis, hematopoiesis, ovulation, menstruation, pregnancy and placentation, or any other disease or condition in which invasion or angiogenesis is pathogenic or undesired.

More recently, it has become apparent that angiogenesis inhibitors may play a role in preventing inflammatory angiogenesis and gliosis following traumatic spinal cord injury, thereby promoting the reestablishment of neuronal connectivity (Wamil, A.W. *et al.*, *Proc. Nat'l. Acad. Sci. USA* 95:13188-13193 (1998)). Therefore, the compositions

of the present invention are administered as soon as possible after traumatic spinal cord injury and for several days up to about two weeks thereafter to inhibit the angiogenesis and gliosis that would sterically prevent reestablishment of neuronal connectivity. The treatment reduces the area of damage at the site of spinal cord injury and facilitates
5 regeneration of neuronal function and thereby prevents paralysis. The compounds of the invention are expected also to protect axons from Wallerian degeneration, reverse aminobutyrate-mediated depolarization (occurring in traumatized neurons), and improve recovery of neuronal conductivity of isolated central nervous system cells and tissue in culture.

10 Having now generally described the invention, the same will be more readily understood through reference to the following examples which are provided by way of illustration, and are not intended to be limiting of the present invention, unless specified.

EXAMPLE I

Binding of the Peptides to HeLa Cells

The compounds of the invention are tested for their binding to uPAR by measuring their ability to inhibit the binding of [125 I]DFP-uPA (catalytically inactivated high molecular weight uPA) to uPAR expressed by HeLa (human cervical carcinoma) cells. Cells (about 5×10^4 /well) are plated (in MEM with Earle's salts/10% FBS +
20 antibiotics) in 24-well plates, then incubated in a humid 5% CO₂ atmosphere until the cells reach 70% confluence. Catalytically inactivated high molecular weight uPA (DFP-uPA) is radioiodinated using Iodo-gen® (Pierce) to a specific activity of about 250,000 cpm/mg. The cell-containing plates are then chilled on ice and the cells washed twice (5 minutes each) with cold PBS/ 0.05% Tween-80. Test compounds are serially diluted in
25 cold PBS/ 0.1 % BSA/ 0.01% Tween-80 and added to each well to a final volume of 0.3mL 10 minutes prior to the addition of the [125 I]DFP-uPA. Each well then receives 9500 cpm of [125 I]DFP-uPA at a final concentration of 0.2 nM. The plates are incubated at 4°C for 2 hrs, after which time the cells are washed 3x (5 minutes each) with cold PBS/ 0.05% Tween-80. NaOH (1N; 0.5 mL) is added to each well to lyse the cells, the
30 contents of each well are aspirated and the total counts in each well determined using a gamma counter. Each compound is tested in triplicate and the results are expressed as a percentage of the total radioactivity measured in wells containing [125 I]DFP-uPA alone, which is taken to represent maximum (100 %) binding.

The peptides of this invention inhibit the binding of [125 I]DFP-uPA with an IC₅₀
35 of ranging between about 0.1 nM to 2 μ M. Modification of the peptides with

diagnostic of therapeutic labels does not significantly affect their measured affinity constants.

EXAMPLE II

Oregon Green-labeled-Peptide Targeting to Tumor Cells

5 The ability of Oregon Green-labeled-peptides to localize to tumor cells is assessed by fluorescence microscopy. Tumor cells are cultured for 24 hours on coverslips (in a 6-well plate). Oregon Green-labeled peptides (1 μ M) are added to each well and allowed to incubate in the presence of the cells. The coverslips are removed from the wells and washed twice (5 minutes per wash) with PBS in a second 6-well
10 plate. The coverslips are mounted onto slides and the fluorescence observed using a fluorescent microscope. Digitized images are captured using a video camera and NIH image. Binding of Oregon Green-labeled-peptides is observed on tumor cells but is generally absent or very weak on normal cells.

EXAMPLE III

Imaging of Tumors *in vivo* by uPAR-targeting Ligand

15 MDA-MB-231 human breast cancer cells (1×10^6) are inoculated s.c. into the flank of female Balb/c (*nu/nu*) mice. Tumors are allowed to grow for varying lengths of time. Labeled uPAR targeting peptides (50 μ g/injection) are injected i.p. and the ability of these peptides to localize to and detect the tumors is assessed at various times post-
20 injection using the appropriate imaging technique.

EXAMPLE IV

Detection of Experimental Metastasis of Tumor Cells *In Vivo* by uPA Peptides

 The peptides described above are also tested *in vivo* in a model of human tumor metastasis in nude mice. PC-3 cells are inoculated into nude mice i.v. at doses of 1×10^6 cells
25 per mouse. These mice are administered the imaging agent as above. The localization of the labeled peptide is imaged using the appropriate detection technique.

 In one study, PC-3 cells are transfected with the gene encoding the enzyme chloramphenicol acetyl-transferase (CAT). At termination of the study, the animals are euthanized and the tumor marker probe CAT is assayed in regional lymph nodes, femurs, lungs,
30 and brain. The localization of the labeled peptide is compared to the actual presence and amount of tumor in a given tissue or site.

 The references cited above are all incorporated by reference herein, whether specifically incorporated or not.

WHAT IS CLAIMED IS:

1. A uPAR-targeting protein or peptide which
 - (a) is diagnostically or therapeutically labeled;
 - (b) comprises at least 38 amino acid residues;
 - 5 (c) includes residues 13-30 of the uPAR-binding site of uPA;
 - (d) competes with labeled DFP-uPA for binding to a cell or molecule that has a binding site for uPA, and has an IC_{50} value of about 10 nM or less; and
 - (e) is not a fusion protein wherein the uPA peptide is fused to another non-uPA protein or peptide.
- 10 2. The protein or peptide of claim 1 selected from the group consisting of:
 - (a) uPA;
 - (b) scuPA;
 - (c) tcuPA;
 - (d) an N-terminal fragment of uPA, residues 1-135;
 - 15 (e) an N-terminal fragment of uPA, residues 1-143;
 - (f) an N-terminal fragment of uPA, residues 1-43; and
 - (g) an N-terminal fragment of uPA, residues 4-43.
- 20 3. A diagnostically useful uPAR-targeting composition comprising:
 - (a) the protein or peptide of claim 1 or 2 which is diagnostically labeled with a detectable label; and
 - (b) a diagnostically acceptable carrier.
- 25 4. The protein, peptide or composition of any of claims 1-3 wherein said label is bound to the protein or peptide through one or more diethylenetriaminepenta-acetic acid residues that are coupled to the protein or peptide.
- 25 5. The composition of any of claims 1 - 3 wherein the detectable label is a radionuclide, a PET-imageable agent, an MRI-imageable agent, a fluorescer, a fluorogen, a chromophore, a chromogen, a phosphorescer, a chemiluminescer or a bioluminescer.
- 30 6. The composition of claim 4 wherein the detectable label is a radionuclide, a PET-imageable agent, an MRI-imageable agent, a fluorescer, a fluorogen, a chromophore and a chromogen.

7. The composition of any of claims 3-6 wherein the detectable label is a radionuclide selected from the group consisting of ^3H , ^{14}C , ^{35}S , ^{67}Ga , ^{68}Ga , ^{72}As , ^{89}Zr , ^{97}Ru , ^{99}Tc , ^{111}In , ^{123}I , ^{125}I , ^{131}I , ^{169}Yb and ^{201}Tl .

8. The composition of any of claims 3-6, wherein the detectable label is a fluorescer or fluorogen selected from the group consisting of fluorescein, rhodamine, dansyl, phycoerythrin, phycocyanin, allophycocyanin, *o*-phthaldehyde, fluorescamine, a fluorescein derivative, Oregon Green, Rhodamine Green, Rhodol Green or Texas Red.

9. The composition of any of claims 3-6 wherein the detectable label is an MRI-imageable agent.

10. 10. The protein, peptide or composition of claim 9 wherein said imageable agent is gadolinium.

11. A uPA active site-targeting compound that covalently modifies the active site of tPA or a fragment or subunit thereof, which fragment or subunit retains (i) the uPA enzymatic endosite and (ii) a uPAR-binding epitope, said compound including a

- 15 (a) detectable label;
(b) a therapeutic moiety; or
(c) a chelator that is optionally bound to a detectable label or a therapeutic moiety;

wherein the compound localizes said chelator, detectable label or therapeutic moiety to the uPA active site.

12. The compound of claim 11 which is an affinity label or a uPA-activated irreversible inhibitor.

13. The compound of claim 12 wherein said affinity label includes the alkylating group chloromethylketone (CMK).

25 14. The compound of claim 11-13 having a structure characterized by the general formula:


(Label)-(Xaa)₂₋₆-(Lys,Arg)-(alkylating group);
(Therapeutic moiety)-(Xaa)₂₋₆-(Lys,Arg)-(alkylating group);
(Chelator_(empty))-(Xaa)₂₋₆-(Lys,Arg)-(alkylating group); or
30 (Label-Chelator)-(Xaa)₂₋₆-(Lys,Arg)-(alkylating group).

wherein Xaa is any amino acid and the label is a detectable label.

15. The compound of claim 14 wherein (Xaa)₂₋₆ is Glu-Gly.
16. The compound of claim 14 having the general formula:
(Chelator_(empty))-Glu-Gly-Arg-CMK; or
(Label-Chelator)-Glu-Gly-Arg-CMK.
- 5 17. A molecule comprising the uPA active site targeting compound of any of claims 11-16, bonded covalently to uPA, tcuPA or a fragment or subunit thereof, which fragment or subunit retains (i) the uPA enzymatic endosite and (ii) a uPAR-binding epitope.
- 10 18. The molecule of claim 17 which has the general formula:
(Chelator_(empty))-(Xaa)₂₋₆-(Lys,Arg)-(alkylating group)-uPA; or
(Label-Chelator)-(Xaa)₂₋₆-(Lys,Arg)-(alkylating group)-uPA.
- 15 19. A uPA active site-targeting peptide compound that binds to the endosite and one or more exosites of (i) tcuPA or (ii) a fragment or subunit of tcuPA, which fragment or subunit retains the uPA (1) enzymatic endosite and (2) a uPAR-binding epitope, such that said peptide compound covalently modifies the endosite; said peptide compound including
- (a) a detectable label,
(b) a therapeutic moiety, or
(c) a chelator that is optionally bound to a detectable label or a therapeutic moiety;
- 20 wherein the peptide compound localizes said chelator, detectable label or therapeutic moiety to the uPA active site.
20. The peptide compound of claim 19 that has a structure defined by a general formula selected from the group consisting of:
- 25 (Label)-(Peptide Z)-(Xaa)₂₋₆-(Lys,Arg)-(alkylating group);
(Therapeutic Moiety)-(Peptide Z)-(Xaa)₂₋₆-(Lys,Arg)-(alkylating group),
(Chelator_(empty))-(Peptide Z)-(Xaa)₂₋₆-(Lys,Arg)-(alkylating group); and
(Label-Chelator)-(Peptide Z)-(Xaa)₂₋₆-(Lys,Arg)-(alkylating group),
wherein, in the formula, the Label is a detectable label, Xaa is any amino acid and Peptide Z
30 is any peptide that binds to a surface exosite of uPA.

21. A molecule comprising the peptide compound of any of claims 19-21 bonded to the uPA endosite and to one or more exosites of
- (a) uPA,
 - (b) tcuPA or
 - 5 (c) a fragment or subunit of tcuPA that retain (i) the uPA enzymatic endosite and (ii) a uPAR-binding epitope.
22. A method for detecting the presence of uPAR (i) on the surface of a cell, (ii) in a tissue, (iii) in an organ or (iv) in a biological sample, which cell, tissue, organ or sample is suspected of expressing uPAR due to a pathological state, comprising the steps of:
- 10 (a) contacting the cell, tissue, organ or sample with the molecule or composition of claim 1-8 or 11-21;
 - (b) detecting the presence of the label associated with the cell, tissue, organ or sample.
23. The method of claim 22, wherein the contacting and the detecting are *in vitro*.
- 15 24. The method of claim 22 wherein the contacting is *in vivo* and the detecting is *in vitro*.
25. The method of claim 22, wherein the contacting and the detecting are *in vivo*.
- 20 26. The method of claim 25, wherein the detectable label is an MRI-imageable agent and the detecting is by MRI.
27. The method of claim 26, wherein the imageable agent is gadolinium.
28. A diagnostic or therapeutic uPAR-targeting pharmaceutical composition comprising:
- 25 (a) an effective amount of the protein or peptide of any of claims 1-4, the compound of any of claims 11-16, the molecule of claim 17 or 18, or a peptide of any of claims 19 or 20, to which is bound directly or indirectly a therapeutically active moiety or a detectable label; and
 - (b) a pharmaceutically acceptable carrier.

29. The pharmaceutical composition of claim 28 in a form suitable for injection.
- 30.. The therapeutic pharmaceutical composition of claim 28 wherein the therapeutically active moiety is a radionuclide.
- 5 31. The therapeutic pharmaceutical composition of claim 29, wherein the radionuclide is selected from the group consisting of ^{47}Sc , ^{67}Cu , ^{90}Y , ^{109}Pd , ^{125}I , ^{131}I , ^{186}Re , ^{188}Re , ^{199}Au , ^{211}At , ^{212}Pb and ^{217}Bi .
- 10 32. A method for inhibiting cell migration, cell invasion, cell proliferation or angiogenesis, or for inducing apoptosis, comprising contacting cells associated with undesired cell migration, invasion, proliferation or angiogenesis with an effective amount of a therapeutic pharmaceutical composition according to any of claims 28-31.
33. A method for inhibiting the invasiveness of tumor cells comprising contacting the cells with an effective amount of a therapeutic pharmaceutical composition according to any of claims 28-31.
- 15 34. A method for treating a subject having a disease or condition associated with undesired cell migration, invasion, proliferation, or angiogenesis, comprising administering to the subject an effective amount of a pharmaceutical composition according to any of claims 28-31.



(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
12 April 2001 (12.04.2001)

PCT

(10) International Publication Number
WO 01/25410 A3

(51) International Patent Classification⁷: C12N 9/72,
C07K 14/81, 5/00, C12Q 1/37, G01N 33/68, A61K 38/49,
38/04, 38/57

(74) Agent: LIVNAT, Shmuel; Venable, P.O. Box 34385,
Washington, DC 20043-9998 (US).

(21) International Application Number: PCT/US00/26502

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ,
DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR,
HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR,
LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ,
NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM,
TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW.

(22) International Filing Date:
27 September 2000 (27.09.2000)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/157,012 1 October 1999 (01.10.1999) US

(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,
IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG,
CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

(71) Applicant: ANGSTROM PHARMACEUTICALS,
INC. [US/US]; Suite 105, 6310 Nancy Ridge Drive, San
Diego, CA 92121 (US).


Published:
— with international search report

(72) Inventors: MAZAR, Andrew, P.; 8650-4 Villa La Jolla
Drive, La Jolla, CA 92037 (US). JONES, Terence, R.;
4915 Quincy Street, San Diego, CA 92109 (US).

(88) Date of publication of the international search report:
10 January 2002

[Continued on next page]

(54) Title: DIAGNOSTIC PROBES AND THERAPEUTICS TARGETING uPA AND uPAR



Urokinase Plasminogen Activator (uPA)

(57) Abstract: A uPAR-targeting protein or peptide is diagnostically or therapeutically labeled and used in methods of diagnosis of therapy. The labeled protein or peptide preferably has the following properties: it comprises at least 38 amino acid residues, including residues 13-30 of the uPAR-binding site of uPA; competes with labeled DFP-uPA for binding to a cell or molecule that has a binding site for uPA, and has an IC_{50} value of about 10 nM or less; and is not a fusion protein wherein the uPA peptide is fused to another non-uPA protein or peptide. Preferred molecules are uPA, scuPA, tcuPA, an N-terminal fragment of uPA, residues 1-135, an N-terminal fragment of uPA, residues 1-143, an N-terminal fragment of uPA, residues 1-43; or an N-terminal fragment of uPA, residues 4-43. Detectable labels include a radionuclide, a PET-imageable agent, an MRI-imageable agent, a fluoroscer, a fluorogen, a chromophore, a chromogen, a phosphorescer, a chemiluminescer or a bioluminescer. The disclosed methods are used to inhibit cell migration, cell invasion, cell proliferation or angiogenesis, or to induce apoptosis, preferably in the treatment of a subject having a disease or condition associated with undesired cell migration, invasion, proliferation or angiogenesis.



For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

INTERNATIONAL SEARCH REPORT

Intern. Appl. No.

PCT/US 00/26502

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C12N9/72 C07K14/81 C07K5/00 C12Q1/37 G01N33/68
A61K38/49 A61K38/04 A61K38/57

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C12N C07K G01N A61K C12Q

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 99 48509 A (RABBANI SHAFAT A ;HART RICHARD (US)) 30 September 1999 (1999-09-30)	1-10, 22-34
Y	the whole document	16,18-34
X	--- DATABASE CAPLUS 'Online! CHEMICAL ABSTRACTS SERVICE, COLUMBUS, OHIO, US; Acc.no. 1986:605286, 1986 HIGGINS, D.L. ET AL.: "The incorporation of a fluorescent probe into the active sites of one- and two-chain tissue-type plasminogen activator." XP002178460	11-15,17
Y	abstract	16,18-34
	--- -/-	



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *S* document member of the same patent family

Date of the actual completion of the international search

25 September 2001

Date of mailing of the international search report

11.10.01

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Smalt, R

INTERNATIONAL SEARCH REPORT

International Application No
PCT/US 00/26502

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	EP 0 890 638 A (NISSIN FOOD PRODUCTS LTD) 13 January 1999 (1999-01-13) the whole document ---	1-10,28, 29,32-34
A	WO 98 21230 A (HANEY DAVID N ;VARGA JANOS (US); JONES TERENCE R (US); ANGSTROM PH) 22 May 1998 (1998-05-22) the whole document ---	
A	APPELLA E ET AL: "THE RECEPTOR-BINDING SEQUENCE OF UROKINASE" JOURNAL OF BIOLOGICAL CHEMISTRY,US,AMERICAN SOCIETY OF BIOLOGICAL CHEMISTS, BALTIMORE, MD, vol. 262, no. 10, 5 April 1987 (1987-04-05), pages 4437-4440, XP000881270 ISSN: 0021-9258 cited in the application the whole document ---	
A	PLOUG E A: "Photoaffinity labelling of the human receptor for urokinase-type plasminogen activator using a decapeptide antagonist. Evidence for a composite ligand-binding site and a short interdomain separation" BIOCHEMISTRY,US,AMERICAN CHEMICAL SOCIETY. EASTON, PA, vol. 37, no. 11, 17 March 1998 (1998-03-17), pages 3612-3622, XP002124882 ISSN: 0006-2960 the whole document ---	
A	WO 90 12091 A (CANCERFORSKNINGSFONDET AF 1989) 18 October 1990 (1990-10-18) page 5, line 13 - line 15 page 22, line 20 - line 21 page 134, line 28 - line 32 -----	

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US 00/26502

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
see FURTHER INFORMATION sheet PCT/ISA/210
2. ☒ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
see FURTHER INFORMATION sheet PCT/ISA/210
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

see additional sheet

1. ☒ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☒ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

This International Searching Authority found multiple (groups of) inventions in this international application, as follows:

1. Claims: 1-10 completely and claims 22-34 partially

Diagnostically or therapeutically labelled uPAR-targeting proteins comprising residues 13-30 of uPA, and their use in treatment and diagnosis.

2. Claims: 11-21 completely, and claims 22-34 partially

Detectably or therapeutically labelled uPA active site-targeting compounds, and their use in treatment and diagnosis.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box I.1

Although claim 25, claim 22 in as far as it pertains to in vivo use, and claim 24 to the extent that the in vivo contacting forms an essential feature of the method, are directed to a diagnostic method practised on the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.

Although claim 34, and claims 32 and 33 in as far as they pertain to in vivo use, are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.

Continuation of Box I.2

Present claims 11-13, 17, 19 and 21 relate to a compound defined by reference to a desirable characteristic or property, namely its ability to target and covalently modify the enzymatic endosite of uPA. The claims cover all compounds having this characteristic or property, whereas the application provides support within the meaning of Article 6 PCT and/or disclosure within the meaning of Article 5 PCT for only a very limited number of such compounds. In the present case, the claims so lack support, and the application so lacks disclosure, that a meaningful search over the whole of the claimed scope is impossible. Independent of the above reasoning, the claims also lack clarity (Article 6 PCT). An attempt is made to define the product by reference to a result to be achieved. Again, this lack of clarity in the present case is such as to render a meaningful search over the whole of the claimed scope impossible. Consequently, the search has been carried out for those parts of the claims which appear to be clear, supported and disclosed, namely those parts relating to the compounds described and claimed in claims 14 and 20.

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.

INTERNATIONAL SEARCH REPORT

Information on patent family members

Intern. Application No

PCT/US 00/26502

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 9948509	A	30-09-1999	US 6077508 A	20-06-2000
			AU 3116099 A	18-10-1999
			EP 1064005 A1	03-01-2001
			WO 9948509 A1	30-09-1999
EP 0890638	A	13-01-1999	AU 1210797 A	01-08-1997
			EP 0890638 A2	13-01-1999
			WO 9725422 A1	17-07-1997
WO 9821230	A	22-05-1998	US 5942492 A	24-08-1999
			AU 4991297 A	03-06-1998
			EP 0941235 A1	15-09-1999
			JP 2001504819 T	10-04-2001
			WO 9821230 A1	22-05-1998
WO 9012091	A	18-10-1990	AT 197067 T	15-11-2000
			AU 652917 B2	15-09-1994
			AU 5528090 A	05-11-1990
			AU 683951 B2	27-11-1997
			AU 8046394 A	13-04-1995
			CA 2051676 A1	08-10-1990
			DE 69033653 D1	23-11-2000
			DE 69033653 T2	07-06-2001
			DE 467932 T1	14-01-1993
			WO 9012091 A1	18-10-1990
			DK 467932 T3	12-02-2001
			EP 1013285 A2	28-06-2000
			EP 0467932 A1	29-01-1992
			JP 4506148 T	29-10-1992
			US 5519120 A	21-05-1996
			US 5891664 A	06-04-1999
			US 6248712 B1	19-06-2001
			US 6113897 A	05-09-2000